

MULTIMEDIA



UNIVERSITY

STUDENT ID NO

--	--	--	--	--	--	--	--	--	--

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 2, 2017/2018

DEM5028 – ENGINEERING MATHEMATICS 2

(Group: E17)

7 MARCH 2018

9.00 a.m. – 11.00 a.m.

(2 Hours)

---

**INSTRUCTIONS TO STUDENT**

1. This question paper consists of 4 pages (2 pages with 4 questions and 2 pages of appendix.
2. Answer **ALL** questions. All necessary working steps must be shown.
3. Write all your answers in the answer booklet provided.

**QUESTION 1 [25 MARKS]**

- a. Using suitable integration method, evaluate the following integrals.

i)  $\int_0^1 x^2(x^3 + 3)^3 dx$

[6 marks]

ii)  $\int 3xe^{2x} dx$

[7 marks]

- b. Find the volume of the enclosed region bounded by  $y = x - 1$  and  $x^2 + y = 1$  that rotate at  $y = -4$ .

[7 marks]

- c. Find the area of the region bounded by the curves  $y = x^2 + 2$ , and  $x + y = 2$ .

[5 marks]

**QUESTION 2 [25 MARKS]**

- a. Find the solution of the differential equation  $y \frac{dy}{dx} = 2x$ .

[4 marks]

- b. For the differential equation of  $\frac{dy}{dx} + y = e^x$ , find the final solution that satisfy the condition of  $y(0) = 1$ .

[9 marks]

- c. Test either the series of  $\sum_{n=1}^{\infty} \frac{8\sqrt{n}}{n^3}$  is convergence or divergence. (Must state the test used).

[3 marks]

- d. Consider the power series  $\sum_{n=1}^{\infty} \frac{nx^n}{3^n}$

- i) Find the radius of convergence.

[4 marks]

- ii) Find the interval of convergence.

[5 marks]

Continued...

**QUESTION 3 [25 MARKS]**

- a. Consider vector  $\mathbf{a} = \langle -2, 2, -2 \rangle$  and  $\mathbf{b} = \langle 1, -3, -3 \rangle$ , find
- i) vector  $\mathbf{b} \times \mathbf{a}$ . [4 marks]
  - ii) the angle between  $\mathbf{a}$  and  $\mathbf{b}$ . [4 marks]
  - iii)  $\mathbf{b} \cdot (\mathbf{a} + 2\mathbf{b})$  [6 marks]
- b. Find the symmetric and parametric equations of the line that goes through the points  $P(1, 2, 4)$  and  $Q(3, -1, 6)$ . [6 marks]
- c. Find the equation of the plane through the point  $(-2, 8, 10)$  and perpendicular to the line  $x = 1 + t, y = 2t, z = 4 - 3t$ . [5 marks]

**QUESTION 4 [25 MARKS]**

- a. For  $f(x, y, z) = x^3z - 3xy^2 - (yz)^3$ , find all its first partial derivatives. [3 marks]
- b. Given  $f(x, y) = x^2 + \frac{1}{3}y^3 - 2xy - 3y$ .
- i) Find the critical point(s) of the function. [4 marks]
  - ii) Determine whether the critical point(s) is a maximum, minimum or saddle point. [8 marks]
- c. A lamina occupies a region between  $x = -1, x = 1, y = 0$  and  $y = 1$  has a density of  $\rho(x, y) = x^2$ .
- i) Find its mass [2 marks]
  - ii) Find its center of mass [8 marks]

**End of page.**

**APPENDIX I: Formulae****Integration of common functions**

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad \int \frac{1}{x} dx = \ln|x| + C \quad \int e^x dx = e^x + C \quad \int a^x dx = \frac{a^x}{\ln a} + C$$

$$\int \sin x dx = -\cos x + C \quad \int \cos x dx = \sin x + C \quad \int \sec^2 x dx = \tan x + C \quad \int \sec x \tan x dx = \sec x + C$$

$$\int \csc^2 x dx = -\cot x + C \quad \int \csc x \cot x dx = -\csc x + C$$

**Inverse Trigonometry****Pythagorean Identities****Integration by parts**

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C$$

$$\sin^2 x + \cos^2 x = 1$$

$$1 + \cot^2 x = \csc^2 x$$

$$\int u dv = uv - \int v du$$

$$\int \frac{1}{x^2+1} dx = \tan^{-1} x + C$$

$$1 + \tan^2 x = \sec^2 x$$

**Areas Between Curves****Volume by Washer****Volume by Cylindrical Shells**

$$A = \int_a^b [f(x) - g(x)] dx$$

$$V = \int_a^b \pi ([f(x)]^2 - [g(x)]^2) dx$$

$$V = \int_a^b 2\pi x(f(x) - g(x)) dx$$

$$A = \int_c^d [w(y) - v(y)] dy$$

$$V = \int_c^d \pi ([w(y)]^2 - [v(y)]^2) dy$$

$$V = \int_c^d 2\pi y(w(y) - v(y)) dy$$

**Linear Differential Equations:**

$$\frac{dy}{dx} + p(x)y = q(x); \mu y = \int \mu q(x) dx \Rightarrow y = \frac{1}{\mu} \int \mu q(x) dx, \quad \text{where } \mu = e^{\int p(x) dx}$$

<b>Divergence Test</b>	If $\lim_{n \rightarrow \infty} a_n \neq 0$ , then $\sum a_n$ diverges.
<b>p-series</b>	The p-series $\sum_{n=1}^{\infty} \frac{1}{n^p}$ is convergent if $p > 1$ and divergent if $p \leq 1$ .
<b>Limit Comparison Test</b>	Let $\sum a_n$ and $\sum b_n$ be series with positive terms such that $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$ If $0 < c < \infty$ , then both series converge or both diverge.
<b>Alternating Series Test</b>	If the alternating series $\sum_{n=1}^{\infty} (-1)^{n-1} b_n = b_1 - b_2 + b_3 - b_4 + \dots \quad b_n > 0$ Satisfies : i. $b_{n+1} \leq b_n$ for all $n$ ii. $\lim_{n \rightarrow \infty} b_n = 0$ then the series is convergent
<b>Ratio Test</b>	Let $\sum a_n$ be a series with nonzero terms such that $L = \lim_{n \rightarrow \infty} \frac{ a_{n+1} }{ a_n }$ a. Series converges absolutely if $L < 1$ b. Series diverges if $L > 1$ or $L = \infty$ c. No conclusion if $L = 1$

Quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Vector

The length of the vector  $\mathbf{a} = \langle a_1, a_2, a_3 \rangle$  is  $|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$ .

If  $\theta$  is the angle between the vector  $\mathbf{a}$  and  $\mathbf{b}$ , then  $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}||\mathbf{b}|\cos\theta$  &  $|\mathbf{a} \times \mathbf{b}| = |\mathbf{a}||\mathbf{b}|\sin\theta$

Cross Product

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \begin{vmatrix} a_2 & a_3 \\ b_2 & b_3 \end{vmatrix} \mathbf{i} - \begin{vmatrix} a_1 & a_3 \\ b_1 & b_3 \end{vmatrix} \mathbf{j} + \begin{vmatrix} a_1 & a_2 \\ b_1 & b_2 \end{vmatrix} \mathbf{k}$$

Equation of Line

Vector equation:  $\mathbf{r} = \mathbf{r}_0 + t\mathbf{v}$

Parametric equation:  $x = x_0 + at, y = y_0 + bt, z = z_0 + ct$

Equation of Plane

$$\langle a, b, c \rangle \cdot \langle x - x_0, y - y_0, z - z_0 \rangle = 0$$

The Chain Rule

Suppose that  $z = f(x, y)$ , where  $x = g(t)$  and  $y = h(t) \Rightarrow \frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$

Second Derivatives Test

Suppose that  $f_x(a, b) = 0$  and  $f_y(a, b) = 0$  [that is,  $(a, b)$  is a critical point of  $f$ ]. Let

$$D = D(a, b) = f_{xx}(a, b)f_{yy}(a, b) - [f_{xy}(a, b)]^2$$

- If  $D > 0$  and  $f_{xx}(a, b) > 0$ , then  $f(a, b)$  is a local minimum.
- If  $D > 0$  and  $f_{xx}(a, b) < 0$ , then  $f(a, b)$  is a local maximum.
- If  $D < 0$ , then  $f(a, b)$  is a saddle point.

Moments and Centers of Mass

The moment about the x-axis:

$$M_x = \iint_D y\rho(x, y)dA$$

The moment about the y-axis:

$$M_y = \iint_D x\rho(x, y)dA$$

The coordinates  $(\bar{x}, \bar{y})$  of the center of mass:

$$\bar{x} = \frac{M_y}{m} = \frac{1}{m} \iint_D x\rho(x, y)dA \quad \bar{y} = \frac{M_x}{m} = \frac{1}{m} \iint_D y\rho(x, y)dA \quad \text{Where the mass: } m = \iint_D \rho(x, y)dA$$

Triple Integrals:  $\iiint_B f(x, y, z)dV = \int_a^s \int_c^d \int_a^b f(x, y, z)dx dy dz$